1. Introduction ..... 4
1.1 Context ..... 4
1.2 Overview of Methodology Document ..... 5
2. Food Losses ..... 7
2.1 Methodology ..... 10
2.1.1 Avoided food losses ..... 10
2.1.2 Exploring different approaches ..... 10
2.1.3 Tool calculations ..... 13
2.2 Benchmark food loss rates ..... 16
2.2.1 Data sources ..... 16
3. Greenhouse Gas Emissions ..... 20
3.1 Production ..... 20
3.1.1 Crops ..... 20
3.1.2 Animal proteins ..... 31
3.2 Emissions due to processes upstream of losses ..... 34
3.2.1 Transport (pre and post processing) ..... 34
3.2.2 Storage (pre and post processing) ..... 36
3.2.3 Processing ..... 37
3.2.4 Retail ..... 38
3.3 Landfill ..... 40
4. References ..... 42
ANNEX ..... 44
Tool Decision Process ..... 45
VBA code ..... 47
Alternative approaches considered ..... 48

The FAO estimates that approximately one third of the food produced in the world is lost or wasted. Given that the world's population is expected to grow, one effective way to meet the demand for more food is to reduce the significant food loss and waste[1] occurring in many parts of the world.

The IFCs activities are addressing the challenge of food loss in many parts of the developing world by making investments to improve storage, transportation, cold chains, and management of food.

The IFC, supported by the Carbon Trust, developed the Food Loss calculator to enable their staff and potentially their clients to quickly and easily estimate the GHG emissions associated with investing in projects that help to reduce food losses. The tool calculates the cumulative GHG emissions associated with preventing food loss and waste at either the farm, transportation, storage, processing, retail or landfill decomposition stages of the value chain. The calculator was designed as a simple tool that IFC staff could easily use to quantify the GHG mitigation benefit derived from supply-chain investments across countries and food types.

As of 2022, the tool covers 50 crops and animal protein products across 117 countries in the following regions:

- East Asia and the Pacific
- Europe and Central Asia
- Latin America and the Caribbean
- Middle East and North Africa
- South Asia
- Sub-Saharan Africa

The tool enables the user to select an agricultural commodity on which to focus the analysis. By providing the

factors for 50 crops and edible livestock products for 117 countries, which are used as default data when no input is provided by the user.

At the same time, it allows the user to use their own inputs to calculate the food loss rate before and after the project, use their own commodity emission factor or provide additional input for increased granularity of results.

It is important to recognise that there is limited quantitative data available regarding food loss rates or associated GHG emissions for different crops and countries. To provide an estimate when user data is not available, the tool utilises FAO food loss and GHG databases to calculate food loss and GHG data points for different crop and country combinations.

This methodology outlines the assumptions used to estimate these food loss rates and GHG emissions values from

: Overview of FAOSTden7.75

The notion of food losses utilised in this work is defined as follows:[2]

Source: SDG 12.3.1: Global Food Loss, p04 re 20tMbheg8ood supply chain

Source: FAO, The State of Food and Agriculture (2019)

The scope of this work includes:

- On-farm post-harvest/slaughter operations (including milking) ;
- Transport (pre- and post-processing);
- Storage (pre- and post-processing, in on-farm or off-farm dedicated facility) .
- Processing
- Retail

Pre-harvest losses, occurring after the commodities are mature but are not harvested, have been excluded from this analysis as they are usually due to economic or environmental events that are outside the control of the individual farmer.

Interpretation of the data included in the FAO Food Loss and Waste database is based on the Global Food Loss Index methodology.[3] The methodology documentation provides some guidelines on how food loss data should be collected and can therefore be used to understand how loss rates within the FAO data set should be interpreted when constructing the baseline loss rate in the tool. While some of the information are relatively straightforward (e.g. , country and commodities), other require some level of interpretation.

The table below is taken from the FAO methodology[4] and represents the sources of information used to define food loss rates at different stages of the value chain.

[3] SDG 12.3.1: Global Food Loss

| Stage | Tool | Notes |
|---|---|---|
| Production<br>(post-harvest losses) | • Sample survey (smallholder farms)<br>• Complete enumeration (large commercial farms) | May cover on-farm storage and on-farm transportation |
| Storage | • Sample survey (smallholder farms)<br>• Inventory data (large storage facilities) | Can include controlled experiments for various length and storage conditions |
| Transport | • Sample survey of trucks (or other transport modes) at destination | Measurements of a product sample at destination |
| Processing | • Agreement with the private sector or through the producer associations.<br>• Company's accounting records.<br>• Complete enumeration or experimental design.<br>• Additional data can come from existing National Industry Processing questionnaires to ascertain | |

3) Estimating food losses that would occur while achieving post-project gross productions levels with the pre-project

| | | |
|---|---|---|
| Net production | | |
| Loss rate | 20% | 10% |
| Food losses | 33.75 | 15 |

The avoided food losses for each approach are reported in the following table:

| | |
|---|---|
| Approach (0): Actual | 5 |
| Approach (1): Pre-project gross production at post-project conditions | 10 |
| Approach (2): Pre-project net production at post-project conditions | 11.11 |
| Approach (3): Post-project gross production at pre-project conditions | 15 |
| Approach (4): Post-project net production at pre-project conditions | 18.75 |

Approach (1) and (2), based on the food losses avoided while achieving the same production

that matches demand in a given market. This means that any additional net production delivered above the (optimal) level achieved by the project would decrease overall efficiency, because the market might not have enough demand to absorb the additional supply. Since approach (3) measures losses resulting from equal levels of gross production, a decrease in the food loss rate causes an increase in net production and results in sub-optimal net production levels.

On the contrary, approach (4) holds for all of our assumptions. It assumes that the net production level achieved by the project and by the baseline are consistent, and the level ET 2 842.04 20



On3 the contrary

characteristics , but the tool can provide a baseline level based on national or regional averages where this is not possible (see section 2.2);

- The production level (in kg or tonnes) observed or expected after the project is implemented, either gross ( ) or net ( ) of food losses. This value must be provided by the user.

Baseline gross production ( ) is defined as the production level that would have been necessary to provide the same amount of net production delivered by the project had the food loss rate not been improved:

Food losses in the baseline scenario ( ) are then calculated as the product of the baseline gross production level ( ) and the food loss rate observed before the project is implemented ( ):

By definition, the net production is the same in both the baseline and project scenarios, so that we have the following equivalence:

where is the amount of food losses observed or expected after the project is implemented:

Finally, food losses avoided by the project are calculated as the difference between baseline and project food losses:

The methodology is illustrated in

Figure 2: Avoided food losses.

animals on-farm are not considered as these are modelled in the emission factors as

The list of commodities included in the FAO database is extensive and not consistent with the commodity nomenclature used within the scope of Food Loss Tool. We therefore needed to map the FAO's data and name conventions to the tool in order to adopt a consistent nomenclature. In some cases, the FAO commodity was too detailed for the purposes of this tool (for e.g., specifying a particular variety of the commodity), while in other cases, differences are simply due to different wording ( e.g., commodities reported in the singular form rather than plural, or vice versa). The classification for FAO commodities that were included in the scope of this project is reported in (annex
).

Once all items in the FAO database were matched with the nomenclature indicated for this project, each commodity was associated to a commodity group. The Tool's commodity group was indicated for each commodity in the scope of this project, while the classification for the FAO commodity group was performed by the Carbon Trust. The latter classification is required to match the Tool's commodity with the commodity group used by FAO for their food loss estimates by region (see step 5 of the methodology in section 0). The classification is reported in annex .

Regional classifications also differ among different sources. For example, in some cases, the region's name from the FAO Food Loss database cannot be matched with the World Bank Group's country classification. To solve this issue, the region name reconciled using the UN sub-region via the M49 country code. This classification is reported in annex
.

Recognising the lack of food loss rate data for many combinations of commodities and countries covered by this project, a step

For step 1 to 4, the tool will return the arithmetic average of all available observations within the period of interest. Stage 5 ensures that the tool can always provide an estimate, if no data is available in the dataset. The food loss rates for stage 5 are reported in the Annex.

the following data points:

1) Average of maize loss rates during transport in Angola;
2) Average of maize loss rates during transport in Sub-Saharan Africa;
3) Average of pulses & grain loss rates during transport in Angola;
4) Average of pulses & grain loss rates during transport in Sub-Saharan Africa;
5) FAO estimate for cereal loss rate during transport in Sub-Saharan Africa.

For the transport and storage stages, the food loss rate calculated with the methodology described above is split between the pre- and post-processing stages to avoid potential double counting. The split is calculated by weighting the distance travelled by the commodity (for transport) and the number of days in storage (for storage). As an example, if the average food loss rate for maize at the transport stage is 10%, and the commodity is transported for 400 km at the pre-processing stage and 600 km at the post-processing stage, then the associated food loss rates are equal to 4% and 6%, respectively.

- : Greenhouse Gas (GHG) emissions from crop residues consist of nitrous oxide gas from decomposition of nitrogen in crop residues left on managed soils.
- : Greenhouse gas (GHG) emissions from manure applied to soils consist of nitrous oxide gas from nitrogen additions to managed soils from treated manure.
- : Greenhouse gas (GHG) emissions from rice cultivation consist of methane gas from the anaerobic decomposition of organic matter in p addy fields. Computed at Tier 1 following the Revised 1996 IPCC Guidelines for National GHG Inventories (IPCC,1997); the IPCC 2000 Good Practice Guidance and Uncertainty Management in National GHG Inventories (IPCC, 2000), and the IPCC Background Papers (IPCC, 2002).
- : Greenhouse gas (GHG) emissions from synthetic fertilizers consist of nitrous oxide gas from synthetic nitrogen additions to managed soils.

emission factor" (see section ), 61 were from Clune's research, 6 from Poore, and the rest from the sources below.

- Emission factors for cotton were sourced from WWF data.[21] As India produces more

contribute specifically to rice production emissions, it was decided to only include the “Rice Cultivation” emissions category. Thus, in some cases rice emission calculated with this tool might be underestimated.

Each country’s emission factor for rice was calculated by dividing each country’s rice cultivation emissions by the rice production :

For countries that do not have rice production and cultivation emissions data – either because rice is not cultivated or because data is missing – a regional average is

same reason why the countries were capped to a maximum value (see - "Outliers – rest of the countries").

Based on each country's emission profile a percentage was calculated about the ratio of $CO_2e$ emissions from $CH_4$ and $N_2O$. Therefore, all of the crops for the same country share the same percentage when calculating $CO_2e$ emissions. For example, if Apples from Argentina have an emission factor of 0.135 $tCO_2e/t$ and based on Argentina's emissions 98% of $CO_2e$ comes from $N_2O$ and 2% comes from $CH_4$. Then it would be 0.132 $tCO_2e/t$ from $N_2O$ and 0.003 $tCO_2e/t$ from $CH_4$.

In some cases, countries have an "Average emission factor per crop" double or even triple than the rest of the countries. The upper bound of the acceptable range is set using Clune/Poore's weighted average maximum value, calculated as the highest possible "Average emission factor per crop" that could be calculated by using the highest global LCA value included in the Clune/Poore papers. If the country is still higher than that, then it means that there must be an error in the reporting, either under-reporting the crop production or overreporting the country emissions. For the sake of having a conservative estimate using this tool, it was necessary to scale down the "Average emission factor per crop" in these countries. This upward capping came in effect in only 3 countries – Botswana, Mongolia, and Trinidad and Tobago.

Additionally, all countries that were below Clune/Poore's minimum of minimum values were also capped. This cut-off point was chosen because this would be the "Average emission factor per crop", if that country only had crop production of the crops with the lowest emission factor and also had calculated its emissions based only on the most conservative global LCAs from Clune/Poore papers. If the country is still lower than that, then it means that there must be an error in the reporting, either under-reporting the emissions or overreporting the country production. For the sake of having an accurate estimate using this tool, it was necessary to scale up the "Average emission factor per crop" in these countries. This downward capping came in effe5B00t.45 6873-5<00520004>-58<004A004500470058>13<0053>-5<00560004>-43<0054>-3<

Regional cradle-to-gate estimates for GHG emission per tonne fertiliser manufactured comes from Brentrup (2018) and is adjusted for emissions per kg N (see sections 3.1.1.5 to 3.1.1.7 for calculating the amount of N) . Since actual emissions from nitrogen manufacture are not




available, the tool uses the average across different ammonium nitrate and urea fertilisers as the ( ). Fertiliser manufacturing is included alongside field emissions to provide a more complete picture of significant emissions that will be avoide d by preventing food loss. Note that the boundary for considering emissions from producing food is different from that used to determine which food losses to measure.

Brentrup et al (Brentrup 2018) used industry data and expert opinion (the authors are predominantly industry experts from Yara, a large fertiliser manufacturer) to generate cradle-to-gate emission factors for a range of 18 common fertilisers in all relevant IFC global regions. They used the online carbon calculator tool provided by Fertilizers Europe for the calculations ("Carbon Footprinting in Fertilizer Production" 2019). We believe this is a good approach because this tool has been verified by DNV and manufacturing company-specific footprint results have been certified by the Carbon Trust for use as emission factors in the Cool Farm Tool ("Cool Farm Tool | An Online Greenhouse Gas, Water, and Biodiversity Calculator" 2019).

Fertilizer manufacturing emissions vary according to the combination of NPK ratio, technology and local energy mix. In particular, different regions vary by the level of $N_2O$ 'scrubbing' technology installed. This technology mitigates (or 'abates') emissions of this high GWP gas and can halve manufacturing emissions - see for example the difference between Western Europe (which includes fertilizer manufactured 100% in the EU-ETS) with East Asia and Pacific (which has relatively little abatement technology installed and high energy emissions). Without project-specific data on fertiliser sourcing (which may be difficult to

obtain) there are some uncertainties, but these are out-weighed by the important contribution of manufacturing to overall emissions.

Data was adapted from the FAO's report on regional livestock emissions (GLEAM).[29] The methodology applies the IPCC tier 2 approach for inventories[30] and GLEAM production data from 2010. LUC (deforestation) from expanded grazing and from feed production were excluded. Emissions from grassland/savannah burning are not included in the GLEAM model for Sub-Saharan Africa, which is inconsistent with FAOSTAT. The tool follows the GLEAMs methodology in order to be conservative (i.e., use the lower emissions estimate available to avoid overestimating emissions avoided).

The FAO data is reported by kg protein, which needs to be converted into kg meat, milk or egg.

For cattle, pig, and poultry, these values were converted from kg carcass protein to per kg carcass and per kg live-weight using the values in table 9.1 in the v2.0 Documentation and table 9.2 in Supplement S1 ("Resources | Global Livestock Environmental Assessment Model (GLEAM) | Food and Agriculture Organization of the United Nations" 2018).

Four emission factors were calculated per region, reflecting predominantly grazed or mixed feeding either with or without feedlot finishing. To include feedlots in the emission factor, the following equation is used:

---

For milk protein conversion, a value of 3.3% protein content was applied (Gerber 2010).

For egg protein conversion, we applied the value of 12.4% protein content in section 9.1.3 of the GLEAM v2.0 Documentation.

[29] Global Livestock Environmental Assessment Model (GLEAM). The GLEAM is a GIS framework that simulates the bio-physical processes and activities along livestock supply chains under a life cycle assessment approach. The aim of GLEAM is to quantify production and use of natural resources in the livestock sector and to identify environmental impacts of livestock in order to contribute to the assessment of adaptation and mitigation scenarios to move towards a more sustainable livestock sector. GLEAM differentiates key stages along livestock supply chains such as feed production, processing and transport; herd dynamics, animal feeding and manure management; and animal products processing and transport. The model captures the specific impacts of each stage, offering a comprehensive and disaggregated picture of livestock production and its use of natural resources. http://www.fao.org/gleam/results/en/

[30] See IPCC - Task Force on National Greenhouse Gas Inventories (2006).

The FAO publications by MacLeod et al (2013) (figure 5) and by Opio et al (2013) (figures 6 and 28) provide a percentage breakdown of average livestock emissions. These percentages were applied to the regional footprints for each Animal Protein emission factor to determine the typical $CH_4$ and $N_2O$ contributions.

The following table summarises the different farming systems for which emission factors were calculated. Full details are available in the GLEAM v2.0 Documentation (section 1.5.2).

| | System | Definition |
|---|---|---|
| Cattle | Grassland | Pastures and rangelands, less than 10 livestock units per hectare. |
| | Mixed | Areas dominated by cropland, >10% feed is crop/by-products |
| | Combination with feedlot | As above but with finishing on specialised units |
| Pigs | | |
| | Backyard system | Mainly subsistence driven or for local markets |
| | Intermediate system | Market-oriented; medium capital input requirements; local feed at least 30% |
| | Industrial system | Market-oriented; high capital requirements; purchased/intensive feed production |
| Poultry | | |
| | Backyard system | Animals producing meat and eggs for the owner and local market, living freely |
| | Layers | Fully market-oriented; high capital input requirements; purchased/intensive feed |
| | Broilers | Fully market-oriented; high capital input requirements; purchased/intensive feed |
| | | |

| | | |
|---|---|---|
| Very poor | 21,393.7 | 2.8% |
| Poor | 20,999.8 | 0.9% |
| Fair | 20,9419 | 0.6% |
| Good | 20,871.1 | 0.3% |
| Very good | 20,829.5 | 0.1% |
| Excellent | 20,818.6 | 0.0% |

The default emissions factor for storage is based upon energy and refrigeration data from the Global Logistics Emissions Council (GLEC) Framework version 2. Table 37 and explanatory text in the GLEC Framework for Logistics Emissions Accounting and Reporting includes survey data from 49 storage sites in Europe and records median ambient (34 sites) and temperature-controlled (15 sites) emissions per tonne. The latter includes emissions from

Users have the option to change this. If the storage of a commodity that is by default ambient but is actually temperature controlled (e.g. to prevent spoilage or for drying), storage emissions (per tonne) are based upon the GLEC temperature controlled emission factor.

The user can alternatively define energy consumption values per tonne. For electricity consumption (kWh/tonne/day) the time is multiplied by the grid emission factor for the country in question (kg $CO_2$e/kWh) and similarly for natural gas (although the emission factor does not vary by country). Default kWh values are provided as a guide only, based upon an assumption that 50% of the GLEC per tonne emissions are generated by electricity and 50% by natural gas. To estimate kWh electricity or gas the following calculation is used: (tonne CO2e / 2) / emission factor per kWh = amount of kWh.

Note – default chilling energy for eggs only includes electricity (no fuels).

Energy data is taken from the broad literature review conducted by Ladha-Sabur et al (2019), which provides energy consumption during processing of a wide variety of foods in various countries. The tool derives average electricity and natural gas usage per commodity group from this data set. Emissions per country and commodity are calculated based upon the country-specific grid and natural gas emission factors.

The approach therefore represents emissions per commodity across a broad approximation of the variety of different processing technologies and products within different countries.

Users may refine the results by entering a processor-specific emission factor or processor-specific electricity and natural gas consumption data.

There are two implications of the processing stage; processing energy and commodity transformation. The latter implies that the outgoing processed product is different from what came in – for example tinned fruit, flour from wheat or carcass meat from live animals. The difference is associated with losses for most commodities and this is modelled by the tool.

The review paper by Ladha-Sabur et al (2019) includes electricity and natural gas consumption data (in MJ) for a wide range of food processing scenarios (at least 100). The foods involv82.55 Tm .55 Tm

Processing of meat and flour generates significant co -products, implying that more than 1 tonne of live animals or wheat are required to make 1 tonne of carcass or flour. The carcass and flour production emission factors take this into account, by applying a conversion factor

| | |
|---|---|
| Vegetables | 100% |

Users can specify the percentage of food losses going to landfill along the supply chain. In addition, the can user can either provide: a specific emission factor per tonne of food losses going to landfill; or additional information on the landfill type and climate zone.

Alternative routes for food not eaten are either considered as alternative use (rather than a loss), such as anaerobic digestion or composting, or difficult to model and so may be represented as landfill by tool users (e.g. crop harvested but left to decompose at field edges).

Barry Callebaut (2015) Chocolate Sustainability Report. https://www.barry-callebaut.com/sites/default/files/2019-01/barry-callebaut-chocolate-sustainability-report-2014-15.pdf

Brentrup, Frank. 2018. “Updated Carbon Footprint Values for Mineral Fertilizer from Different World Regions.” In .

“Carbon Footprinting in Fertilizer Production.” 2019. (blog). 2019. https://www.fertilizerseurope.com/initiatives/carbon-footprint-calculator/

“Cool Farm Tool | An Online Greenhouse Gas, Water, and Biodiversity Calculator.” n.d. Accessed June 26, 2020. https://coolfarmtool.org/

Clune S, Crossin E, Verghese K (2016) Systematic review of greenhouse gas emissions for different fresh food categories, Journal of Cleaner Production. DOI: 10.1016/j.jclepro.2016.04.08nces

Gilchrist, Christie, Fraser & Inglis (2012). Project Report No. 484; The relationship between soil mineral nitrogen, applied nitrogen and yields in Scottish soils. https://projectblue.blob.core.windows.net/media/Default/Research%20Papers/Cereals%20and%20Oilseed/pr484.pdf

GLEC, Global Logistics Emissions Council Framework version 2. https://www.smartfreightcentre.org/en/how-to-implement-items/what-is-glec-framework/58/

Global Livestock Environmental Assessment Model http://www.fao.org/gleam/en/

FL = Food Losses
Tool uses user
Tool uses user
defined

EF = Emission Factor
Does the user..
the commodity
uses and/or how
commodity has?
Crop
No
Crop or animal
desired country
Start
user know
Yes
Selection
Tool uses
direct user
Animal
production type

In this section some of assumptions used for calculating greenhouse gases from crops will be discussed. Here also previous methodologies that were considered will be discussed and why they were not selected in the end.

- I

- All of the above methodologies' results were compared to each other and to the results of the previous "IFC Crop GHG Calculator".[41]
- Each country's emissions were calculated bottom up by adding all the emissions from all crops. Emissions for each target crop were calculated by multiplying

:

This document and the intellectual property, concepts and content contained within it shall n ot be used for any purpose other than that for which it was provided by the Carbon Trust and shall not be reproduced in whole or in part.

Whilst reasonable steps have been taken to ensure that the information contained within this report is correct, the authors, the Carbon Trust, its agents, contractors and sub-contractors give no warranty and make no representation as to its accuracy and accept no liability for any errors or omissions.

The Carbon Trust is a company limited by guarantee and registered in England and Wales under company number 4190230 with its registered office at 4 th Floor Dorset House, Stamford Street, London, SE1 9NT.

© The Carbon Trust 2022. All rights reserved.